


# Climate4you update January 2011



[www.climate4you.com](http://www.climate4you.com)

## January 2011 global surface air temperature overview





1

Air temperature 201101 versus average 1998-2006



Air temperature 201101 versus average 1998-2006



January 2011 surface air temperature compared to the average 1998-2006. Green,yellow-red colours indicate areas with higher temperature than the 1998-2006 average, while blue colours indicate lower than average temperatures. Data source: [Goddard Institute for Space Studies](http://www.giss.nasa.gov) (GISS)

## Comments to the January 2011 global surface air temperature overview

General: This newsletter contains graphs showing a selection of key meteorological variables for January 2011. All temperatures are given in degrees Celsius.

In the above maps showing the geographical pattern of surface air temperatures, the period 1998-2006 is used as reference period. The reason for comparing with this recent period instead of the official WMO 'normal' period 1961-1990, is that the latter period is affected by the relatively cold period 1945-1980. Almost any comparison with such a low average value will therefore appear as high or warm, and it will be difficult to decide if modern surface air temperatures are increasing or decreasing. Comparing with a more recent period overcomes this problem. In addition to this consideration, the recent temperature development suggests that the time window 1998-2006 may roughly represent a global temperature peak. If so, negative temperature anomalies will gradually become more and more widespread as time goes on. However, if positive anomalies instead gradually become more widespread, this reference period only represented a temperature plateau.

In the other diagrams in this newsletter the thin line represents the monthly global average value, and the thick line indicate a simple running average, in most cases a simple moving 37-month average, almost corresponding to three years. The 37-month average is calculated from values covering a range from 18 month before to 18 months after, with equal weight for every month.

The year 1979 has been chosen as starting point in several of the diagrams, as this roughly corresponds to both the beginning of satellite observations and the onset of the late 20<sup>th</sup> century warming period. Several of the records, however, have a much longer history, which may be inspected on [www.Climate4you.com](http://www.Climate4you.com).

Global surface air temperatures January 2011 in general was below the 1998-2006 average, as detailed by several of the temperature diagrams shown below.

---

2

The Northern Hemisphere was characterised by generally low temperatures, but also by high regional contrasts. A zone of below average temperatures extended from China, across Mongolia, southern Siberia, Europe, the North Atlantic, USA, and southern and western Canada, somewhat like that characterising the previous month (December 2010). Above average temperatures characterised northeast Canada, western Greenland, northern and eastern Siberia, and Alaska.

The Southern Hemisphere in general was close to or slightly below average 1998-2006 conditions. Most of South America and Africa south of Equator experienced below average temperatures. Most of Australia had temperatures slightly above average. There were, however, no major warm regions in the Southern Hemisphere in January 2011.


Near Equator temperatures conditions were influenced by the La Nina situation. Relatively low temperatures characterised most of the Equatorial regions in the Pacific and Indian Ocean. The Equatorial Atlantic was close to average conditions. Because of the huge areas represented by these regions at low latitudes, the cooling effect on the global average temperature is now clearly felt (see below).

The Arctic was characterized by huge contrasts as to surface air temperatures. Most areas experienced above average 1998-2006 temperatures, but northwest Canada, eastern Greenland, and the European Arctic had below average temperatures.

In the Antarctic temperature conditions were slightly above the average for 1998-2006. The Antarctic Peninsula, however, had below average temperatures.


All diagrams shown in this newsletter are available for download on [www.climate4you.com](http://www.climate4you.com)

## Lower troposphere temperature from satellites, updated to January 2011




Global monthly average lower troposphere temperature (thin line) since 1979 according to [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37 month average.

3




Global monthly average lower troposphere temperature (thin line) since 1979 according to according to [Remote Sensing Systems](#) (RSS), USA. The thick line is the simple running 37 month average.

## Global surface air temperature, updated to January 2011




Global monthly average surface air temperature (thin line) since 1979 according to according to the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK. The thick line is the simple running 37 month average.

4





Global monthly average surface air temperature (thin line) since 1979 according to according to the [Goddard Institute for Space Studies \(GISS\)](#), at Columbia University, New York City, USA. The thick line is the simple running 37 month average.



Global monthly average surface air temperature since 1979 according to according to the [National Climatic Data Center](http://www.ncdc.noaa.gov) (NCDC), USA. The thick line is the simple running 37 month average.

Some readers have noted that the above temperature estimates display changes when one compare with previous issues of this newsletter, not only for the most recent months, but actually for all months back to the beginning of the record. As an example, the net change of the NCDC record since 17 May 2008 is shown below. By this administrative effort the apparent global temperature increase since 1900 has been enhanced about  $0.1^{\circ}\text{C}$ , or about 14% of the total increase recorded since 1900 by NCDC. The interested reader may find more on this lack of temporal stability on [www.climate4you](http://www.climate4you) (go to: Global Temperature and then Temporal Stability).





6

*Superimposed plot of all five global monthly temperature estimates shown above. As the base period differs for the different temperature estimates, they have all been normalised by comparing to the average value of their initial 120 months (10 years) from January 1979 to December 1988. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of all five temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the above average.*

It should be kept in mind that satellite- and surface-based temperature estimates are derived from different types of measurements, and that comparing them directly as done in the diagram above therefore in principle is problematical. However, as both types of estimate often are discussed together, the above diagram may nevertheless be of interest. In fact, the different types of temperature estimates appear to agree quite well as to the overall temperature variations on a 2-3 year scale, although on a short term scale there may be considerable differences.


All five global temperature estimates presently show stagnation, at least since 2002. There has been no increase in global air temperature since 1998, which was affected by the oceanographic El Niño event. This does not exclude the possibility that global temperatures will begin to increase again later. On the other hand, it also remain a possibility that Earth just now is passing a temperature peak, and that global temperatures will begin to decrease within the coming 5-10 years. Only time will show which of these possibilities is correct.



## Global sea surface temperature, updated to end of January 2011

NOAA/NWS/NCEP/EMC Marine Modeling and Analysis Branch


RTG\_SST Anomaly (0.5 deg X 0.5 deg) for 30 Jan 2011



*Sea surface temperature anomaly at 30 January 2011. Map source: National Centers for Environmental Prediction (NOAA).*


The relative cold surface water now dominating the Equator in the Pacific Ocean represents a La Niña situation and affects the temperature of the atmosphere above. Because of the large surface areas involved (near Equator) this natural cyclic development is at the moment affecting the global atmospheric temperature towards lower temperatures.

However, the significance of any such global cooling should not be over interpreted. Whenever Earth experiences cold La Niña or warm El Niño episodes major heat exchanges takes place between the Pacific Ocean and the atmosphere above, eventually showing up in estimates of the global air temperature. This does not, however, reflect similar changes in the total heat content of the atmosphere-ocean system. In fact, net changes may be small, as it mainly reflects a redistribution of energy. What matters is the overall development when seen over some years.




Global monthly average lower troposphere temperature over oceans (thin line) since 1979 according to [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37 month average.

8




Global monthly average sea surface temperature since 1979 according to University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK. Base period: 1961-1990. The thick line is the simple running 37 month average.






Global monthly average sea surface temperature since 1979 according to the [National Climatic Data Center](#) (NCDC), USA. Base period: 1901-2000. The thick line is the simple running 37 month average.

## Global ocean heat content, updated to December 2010




Global monthly heat content anomaly (GJ/m<sup>2</sup>) in the uppermost 700 m of the oceans since January 1979. Data source: National Oceanographic Data Center(NODC).



Global monthly heat content anomaly (GJ/m<sup>2</sup>) in the uppermost 700 m of the oceans since January 1955. Data source: National Oceanographic Data Center(NODC).

## Arctic and Antarctic lower troposphere temperature, updated to January 2011



Global monthly average lower troposphere temperature since 1979 for the North Pole and South Pole regions, based on satellite observations ([University of Alabama](http://climate4you.org) at Huntsville, USA). The thick line is the simple running 37 month average, nearly corresponding to a running 3 yr average.

## Arctic and Antarctic surface air temperature, updated to December 2010




Diagram showing Arctic monthly surface air temperature anomaly 70-90°N since January 2000, in relation to the WMO reference "normal" period 1961-1990. The thin blue line shows the monthly temperature anomaly, while the thicker red line shows the running 13 month average. Data provided by the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK.




Diagram showing Antarctic monthly surface air temperature anomaly 70-90°S since January 2000, in relation to the WMO reference "normal" period 1961-1990. The thin blue line shows the monthly temperature anomaly, while the thicker red line shows the running 13 month average. Data provided by the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK.




Diagram showing Arctic monthly surface air temperature anomaly 70-90°N since January 1957, in relation to the WMO reference “normal” period 1961-1990. The year 1957 has been chosen as starting year, to ensure easy comparison with the maximum length of the realistic Antarctic temperature record shown below. The thin blue line shows the monthly temperature anomaly, while the thicker red line shows the running 13 month average. Data provided by the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK.

13




Diagram showing Antarctic monthly surface air temperature anomaly 70-90°S since January 1957, in relation to the WMO reference “normal” period 1961-1990. The year 1957 was an international geophysical year, and several meteorological stations were established in the Antarctic because of this. Before 1957, the meteorological coverage of the Antarctic continent is poor. The thin blue line shows the monthly temperature anomaly, while the thicker red line shows the running 13 month average. Data provided by the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK.




Diagram showing Arctic monthly surface air temperature anomaly 70-90°N since January 1900, in relation to the WMO reference “normal” period 1961-1990. The thin blue line shows the monthly temperature anomaly, while the thicker red line shows the running 13 month average. In general, the range of monthly temperature variations decreases throughout the first 30-50 years of the record, reflecting the increasing number of meteorological stations north of 70°N over time. Especially the period from about 1930 saw the establishment of many new Arctic meteorological stations, first in Russia and Siberia, and following the 2nd World War, also in North America. Because of the relatively small number of stations before 1930, details in the early part of the Arctic temperature record should not be over interpreted. The rapid Arctic warming around 1920 is, however, clearly visible, and is also documented by other sources of information. The period since 2000 is warm, about as warm as the period 1930-1940. Data provided by the Hadley Centre for Climate Prediction and Research and the University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK

In general, the Arctic temperature record appears to be less variable than the contemporary Antarctic record, presumably at least partly due to the higher number of meteorological stations north of 70°N, compared to the number of stations south of 70°S.


As data coverage is sparse in the Polar Regions, the procedure of Gillett et al. 2008 has been followed, giving equal weight to data in each 5°x5° grid cell when calculating means, with no weighting by the areas of the grid cells.

#### Literature:

Gillett, N.P., Stone, D.A., Stott, P.A., Nozawa, T., Karpechko, A.Y.U., Hegerl, G.C., Wehner, M.F. and Jones, P.D. 2008. Attribution of polar warming to human influence. *Nature Geoscience* 1, 750-754.




## Arctic and Antarctic sea ice, updated to January 2011



15


Graphs showing monthly Antarctic, Arctic and global sea ice extent since November 1978, according to the [National Snow and Ice data Center](#) (NSIDC).



Graph showing daily Arctic sea ice extent since June 2002, to 17/02 2011, by courtesy of [Japan Aerospace Exploration Agency](#) (JAXA).


PIPS2.0 24hr forecast from 2010021700\_024.dat  
valid for 2010021800

PIPS2.0 24hr forecast from 2011021700\_024.dat  
valid for 2011021800




*Northern hemisphere sea ice thickness on 18 February 2010 (left) and 2011 (right), according to the Naval Oceanographic Office (NAVO). Thickness values are calculated by the Polar Ice Prediction System (PIPS 2.0), based on the Special Sensor Microwave Image (SSM/I) to initialize the calculation. Thickness scale (m) is shown to the right.*

## Global sea level, updated to September 2010




Global monthly sea level since late 1992 according to the Colorado Center for Astrodynamics Research at [University of Colorado at Boulder](http://www.ccar.colorado.edu), USA. The thick line is the simple running 37 observation average, nearly corresponding to a running 3 yr average.

17




Annual change of global sea level since late 1992 according to the Colorado Center for Astrodynamics Research at [University of Colorado at Boulder](http://www.ccar.colorado.edu), USA. The thick line is the simple running 3 yr average.

## Atmospheric CO<sub>2</sub>, updated to January 2011




18




Monthly amount of atmospheric CO<sub>2</sub> (above) and annual growth rate (below; average last 12 months minus average preceding 12 months) of atmospheric CO<sub>2</sub> since 1959, according to data provided by the [Mauna Loa Observatory](#), Hawaii, USA. The thick line is the simple running 37 observation average, nearly corresponding to a running 3 yr average.

## Northern Hemisphere weekly snow cover, updated to late January 2011




Northern hemisphere weekly snow cover since January 2000 according to Rutgers University Global Snow Laboratory. The thin line is the weekly data, and the thick line is the running 53 week average (approximately 1 year).




Northern hemisphere weekly snow cover since October 1966 according to Rutgers University Global Snow Laboratory. The thin line is the weekly data, and the thick line is the running 53 week average (approximately 1 year). The running average is not calculated before 1971 because of some data irregularities in this early period.


## Global surface air temperature and atmospheric CO<sub>2</sub>, updated to January 2011



20







Diagrams showing HadCRUT3, GISS, and NCDC monthly global surface air temperature estimates (blue) and the monthly atmospheric CO<sub>2</sub> content (red) according to the [Mauna Loa Observatory](#), Hawaii. The Mauna Loa data series begins in March 1958, and 1958 has therefore been chosen as starting year for the diagrams. Reconstructions of past atmospheric CO<sub>2</sub> concentrations (before 1958) are not incorporated in this diagram, as such past CO<sub>2</sub> values are derived by other means (ice cores, stomata, or older measurements using different methodology, and therefore are not directly comparable with modern atmospheric measurements). The dotted grey line indicates the approximate linear temperature trend, and the boxes in the lower part of the diagram indicate the relation between atmospheric CO<sub>2</sub> and global surface air temperature, negative or positive.


Most climate models assume the greenhouse gas carbon dioxide CO<sub>2</sub> to influence significantly upon global temperature. Thus, it is relevant to compare the different global temperature records with measurements of atmospheric CO<sub>2</sub>, as shown in the diagrams above. Any comparison, however, should not be made on a monthly or annual basis, but for a longer time period, as other effects (oceanographic, clouds, etc.) may well override the potential influence of CO<sub>2</sub> on short time scales such as just a few years.

It is of cause equally inappropriate to present new meteorological record values, whether daily, monthly or annual, as support for the hypothesis ascribing high importance of atmospheric CO<sub>2</sub> for global temperatures. Any such short-period meteorological record value may well be the result of other phenomena than atmospheric CO<sub>2</sub>.

What exactly defines the critical length of a relevant time period to consider for evaluating the alleged high importance of CO<sub>2</sub> remains elusive, and is still a topic for debate. The critical period length must, however, be inversely proportional to the importance of CO<sub>2</sub> on the global temperature, including feedback effects, such as assumed by most climate models. So if the effect of CO<sub>2</sub> is strong, the length of the critical period is short.

After about 10 years of global temperature increase following global cooling 1940-1978, IPCC was established in 1988. Presumably, several scientists interested in climate then felt intuitively that their empirical and theoretical understanding of climate dynamics was sufficient to conclude about the high importance of CO<sub>2</sub> for global temperature. However, for obtaining public and political support for the CO<sub>2</sub>-hypothesis the 10 year warming period leading up to 1988 in all likelihood was important. Had the global temperature instead been decreasing, public support for the hypothesis would have been difficult to obtain. Adopting this approach as to critical time length, the varying relation (positive or negative) between global temperature and atmospheric CO<sub>2</sub> has been indicated in the lower panels of the three diagrams above.

**1895: Arvid Högbom's geochemical investigations on atmospheric CO<sub>2</sub>**



*The volcano Stomboli, Italy (left). Swedish geologist Arvid Gustav Högbom (centre). Limestone rocks exposed on the island Møn, southeastern Denmark (right). These chalk formations were deposited in a tropical ocean covering Denmark about 65-70 mill. years ago, and were later pushed into their present, elevated position by the North European Ice Sheet, about 20,000 years ago, when Arctic climate conditions prevailed in most of Europe. By this, the above scenic Danish landscape is a geological visualisation of the existence of significant, natural global climate changes.*

Simultaneously with the publication of Svante Arrhenius 1895 paper (see previous newsletter) on the possible role for atmospheric CO<sub>2</sub> as a control on glaciations, the Swedish geologist Arvid Gustav Högbom was working on the geochemistry of carbon. Arrhenius was using the findings of Högbom as an inspiration for describing a possible mechanism for variations in the amount of atmospheric CO<sub>2</sub>. Arrhenius and Högbom were both living in Stockholm; they were friends and colleagues, and of course followed the research findings of each other with a keen interest. Following the acceptance of the glacial hypothesis after the observations of Louis Agassiz in Scotland in 1840, the mechanisms behind the onset of glaciations were becoming a scientific issue of widespread interest.

From Högbom's perspective, neither the use of fossil fuels nor the removal of organic carbon (deforestation) influenced atmospheric CO<sub>2</sub> nearly as much as a suite of different geological processes. The formation of limestone and other carbonates was mentioned as one important mechanism, by which CO<sub>2</sub> is removed from the atmosphere, and the decomposition of silicates, which adds to the atmospheric amount of CO<sub>2</sub>, was another mechanism. Also volcanic activity was emphasized by Högbom as an important agent adding CO<sub>2</sub> to the atmosphere. Actually, Högbom considered volcanoes to be the "chief source of carbonic acid for the

atmosphere" (Fleming 1998). In addition to this, he also mentioned the combustion of carbonaceous meteorites in the atmosphere as a possible, but virtually unknown, source for CO<sub>2</sub>.

Högbom estimated that the current atmospheric concentration of CO<sub>2</sub> was of the same order of magnitude as the amount of carbon fixed in the living organic world. He further concluded that about 25,000 times as much carbonic acid is fixed in sedimentary formations in the form of limestone (see photo above) as is found in the atmosphere. On top of all this was the regulative role of the oceans, the potential importance of which was indicated by the research findings on the high solubility of CO<sub>2</sub> in water by the English chemist William Henry as early as in 1803. As the carbon cycle thus contained several important processes, and because these in general could be considered independent of each other, Högbom stated that, from a geological point of view, a stable level of atmospheric CO<sub>2</sub> was unlikely to persist at any time. In conclusion he therefore specified that the amount of atmospheric CO<sub>2</sub> was likely to have varied considerably over time, as the result of different geological processes (Fleming 1998).

Having the assurance of Högbom that large variations in the atmospheric concentration of CO<sub>2</sub> were quite likely in different geological periods, Svante Arrhenius in 1895 adopted this as foundation for his hypothesis explaining the onset of ice ages and interglacials as the possible result of natural variations of atmospheric CO<sub>2</sub>.

#### References:

Fleming, J.R. 1998. *Historical Perspectives on Climate Change*. Oxford University Press, 194 pp.

All the above diagrams with supplementary information, including links to data sources and previous issues of this newsletter, are available on [www.climate4you.com](http://www.climate4you.com)

Yours sincerely, Ole Humlum (Ole.Humlum@geo.uio.no)

20 February 2011.