

Climate4you update May 2020



Contents:

- Page 2: May 2020 global surface air temperature overview
- Page 3: May 2020 global surface air temperature overview versus May last 10 years
- Page 4: May 2020 global surface air temperature compared to May 2019
- Page 5: Temperature quality class 1: Lower troposphere temperature from satellites
- Page 6: Temperature quality class 2: HadCRUT global surface air temperature
- Page 7: Temperature quality class 3: GISS and NCDC global surface air temperature
- Page 10: Comparing global surface air temperature and satellite-based temperatures
- Page 11: Global air temperature linear trends
- Page 12: Global temperatures: All in one, Quality Class 1, 2 and 3
- Page 14: Global sea surface temperature
- Page 17: Ocean temperature in uppermost 100 m
- Page 19: North Atlantic heat content uppermost 700 m
- Page 20: North Atlantic temperatures 0-800 m depth along 59N, 30-0W
- Page 21: Global ocean temperature 0-1900 m depth summary
- Page 22: Global ocean net temperature change since 2004 at different depths
- Page 23: La Niña and El Niño episodes
- Page 24: Troposphere and stratosphere temperatures from satellites
- Page 25: Zonal lower troposphere temperatures from satellites
- Page 26: Arctic and Antarctic lower troposphere temperatures from satellites
- Page 27: Temperature over land versus over oceans
- Page 28: Arctic and Antarctic surface air temperatures
- Page 31: Arctic and Antarctic sea ice
- Page 35: Sea level in general
- Page 36: Global sea level from satellite altimetry
- Page 37: Global sea level from tide gauges
- Page 38: Snow cover; Northern Hemisphere weekly and seasonal
- Page 40: Atmospheric specific humidity
- Page 41: Atmospheric CO₂
- Page 42: Relation between annual change of atm. CO₂ and La Niña and El Niño episodes
- Page 43: Phase relation between atmospheric CO₂ and global temperature
- Page 44: Global air temperature and atmospheric CO₂
- Page 48: Latest 20-year QC1 global monthly air temperature change
- Page 49: Sunspot activity and QC1 average satellite global air temperature
- Page 50: Climate and history: *1815: Tambora eruption and cold summer in New England 1816*

May 2020 global surface air temperature overview

General: This newsletter contains graphs and diagrams showing a selection of key meteorological variables, if possible updated to the most recent past month. All temperatures are given in degrees Celsius.

In the maps on pages 3-4, showing the geographical pattern of surface air temperature anomalies, the last previous 10 years are used as reference period.

The rationale for comparing with this recent period instead of the official WMO so-called 'normal' period 1961-1990, is that the latter period is affected by the cold period 1945-1980. Most comparisons with this period will inevitably appear as warm, and it will be difficult to decide if modern temperatures are increasing or decreasing. Comparing instead with the last previous 10 years overcomes this problem and clearer displays the modern dynamics of ongoing change. This decadal approach also corresponds well to the typical memory horizon for many people and is now also adopted as reference period by other institutions, e.g. the Danish Meteorological Institute (DMI).

2

In addition, most temperature databases display temporal instability for data before the turn of the century (see, e.g., p. 8). Any comparison with the WMO reference period 1961-1990 will therefore be influenced by such ongoing monthly changes of mainly administrative nature. An unstable value is clearly not suited as reference value. Simply comparing with the last previous 10 years is more useful as reference for modern changes. See also additional reflections on page 47.

The different air temperature records have been divided into three quality classes, QC1, QC2 and QC3, respectively, as described on page 8.

In many diagrams shown in this newsletter the thin line represents the monthly global average value, and the thick line indicate a simple running average, in most cases a simple moving 37-month average, nearly corresponding to a three-year average. The 37-month average is calculated from values covering a range from 18 months before to 18 months after, with equal weight given to all individual months.

The year 1979 has been chosen as starting point in many diagrams, as this roughly corresponds to both the beginning of satellite observations and the onset of the late 20th century warming period. However, several of the data series have a much longer record length, which may be inspected in greater detail on www.climate4you.com.

May 2020 Remote Sensed Surface Temperature


General: For May 2020 AIRS supplied 16200 interpolated surface air data points, obtained from the GISS data portal. According to most surface temperature databases, the May 2020 global average air temperature anomaly was somewhat lower than estimated for the previous month.

The Northern Hemisphere 10-yr temperature anomaly pattern (p.3) was characterised by large regional contrasts, mainly controlled by the dominant jet stream configuration. Canada, eastern USA, N and E Europe, W Russia, India, W China, and SE Siberia all had temperatures below the average for the previous 10 years. The May peak warm anomaly was in western Siberia, much like in April. Ocean wise, part of the North Atlantic, the Greenland- and the Norwegian Sea were relatively cold. The North Pacific and Alaska both was above the 10-yr average. In the Arctic relatively low temperatures dominated in the Canadian and European sectors, while the remaining arctic regions were relatively warm. Over the central Arctic Ocean surface air temperatures were a little above the average.

Near the Equator temperatures were mostly near or slightly above the 10-year average.


The Southern Hemisphere temperatures were generally near or below the average for the previous 10 years. A zone of relatively low surface air temperatures dominated between 20°S and 50°S, affecting ocean as well as land areas, especially Australia and Brazil. In contrast, southern South America was relatively warm. In the Antarctic, regional surface air temperatures were mainly above the 10-year average, although some regions had below average temperatures.

May 2020 global surface air temperature overview versus average May last 10 years




3

Surface air temperature May 2020 versus last 10 yr




Surface air temperature May 2020 versus last 10 yr





May 2020 surface air temperature compared to the average of May over the last 10 years. Green-yellow-red colours indicate areas with higher temperature than the 10-year average, while blue colours indicate lower than average temperatures. Data source: Remote Sensed Surface Temperature Anomaly, AIRS/Aqua L3 Monthly Standard Physical Retrieval 1 degree x 1 degree V006 (<https://airs.jpl.nasa.gov/>), obtained from the GISS data portal.

May 2020 global surface air temperature compared to May 2019





4

Surface air temperature May 2020 versus May 2019




Surface air temperature May 2020 versus May 2019




May 2020 surface air temperature compared to May 2019. Green-yellow-red colours indicate regions where the present month was warmer than last year, while blue colours indicate regions where the present month was cooler than one year ago. Variations in monthly temperature from one year to the next has no tangible climatic importance but may nevertheless be interesting to study. Data source: Remote Sensed Surface Temperature Anomaly, AIRS/Aqua L3 Monthly Standard Physical Retrieval 1 degree x 1 degree V006 (<https://airs.jpl.nasa.gov/>), obtained from the GISS data portal.

Temperature quality class 1: Lower troposphere temperature from satellites, updated to May 2020




Global monthly average lower troposphere temperature (thin line) since 1979 according to [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37-month average.

5




Global monthly average lower troposphere temperature (thin line) since 1979 according to according to [Remote Sensing Systems](#) (RSS), USA. The thick line is the simple running 37-month average.

Temperature quality class 2: HadCRUT global surface air temperature, updated to April 2020




Temperature quality class 3: GISS and NCDC global surface air temperature, updated to May 2020



7

Global monthly average surface air temperature (thin line) since 1979 according to according to the [Goddard Institute for Space Studies \(GISS\)](#), at Columbia University, New York City, USA, using ERSST_v4 ocean surface temperatures. The thick line is the simple running 37-month average.



Global monthly average surface air temperature since 1979 according to according to the [National Climatic Data Center \(NCDC\)](#), USA. The thick line is the simple running 37-month average.

A note on data record stability and -quality:

The temperature diagrams shown above all have 1979 as starting year. This roughly marks the beginning of the recent episode of global warming, after termination of the previous episode of global cooling from about 1940. In addition, the year 1979 also represents the starting date for the satellite-based global temperature estimates (UAH and RSS). For the three surface air temperature records (HadCRUT, NCDC and GISS), they begin much earlier (in 1850 and 1880, respectively), as can be inspected on www.climate4you.com.

For all three surface air temperature records, but especially NCDC and GISS, administrative changes to anomaly values are quite often introduced, even affecting observations many years back in time. Some changes may be due to the delayed addition of new station data or change of station location, while others probably have their origin in changes of the technique adopted to calculate average values. It is clearly impossible to evaluate the validity of such administrative changes for the outside user of these records; it is only possible to note that such changes quite often are introduced (see example diagram next page).

In addition, the three surface records represent a blend of sea surface data collected by moving ships or by other means, plus data from land stations of partly unknown quality and unknown degree of representativeness for their region. Many of the land stations also has been moved geographically during their period of operation, their instrumentation have been changed, and they are influenced by changes in their near surroundings (vegetation, buildings, etc.).

The satellite temperature records also have their problems, but these are generally of a more technical nature and therefore better correctable. In addition, the temperature sampling by satellites is more regular and complete on a global basis than that represented by the surface records. It is also

important that the sensors on satellites measure temperature directly by emitted radiation, while most modern surface temperature measurements are indirect, using electronic resistance.

Everybody interested in climate science should gratefully acknowledge the big efforts put into maintaining the different temperature databases referred to in the present newsletter. At the same time, however, it is also important to realise that all temperature records cannot be of equal scientific quality. The simple fact that they to some degree differ shows that they cannot all be correct.

On this background, and for practical reasons, Climate4you therefore operates with three quality classes (1-3) for global temperature records, with 1 representing the highest quality level:

Quality class 1: The satellite records (UAH and RSS).

Quality class 2: The HadCRUT surface record.

Quality class 3: The NCDC and GISS surface records.

The main reason for discriminating between the three surface records is the following:

While both NCDC and GISS often experience quite large administrative changes (see example on p.8), and therefore essentially are unstable temperature records, the changes introduced to HadCRUT are fewer and smaller. For obvious reasons, as the past does not change, any record undergoing continuing changes cannot describe the past correctly all the time. Frequent and large corrections in a database of cause signal a fundamental doubt about what is likely to represent the correct values.

You can find more on the issue of lack of temporal stability on www.climate4you.com (go to: *Global Temperature*, and then proceed to *Temporal Stability*).





Diagram showing the adjustments made since May 2008 by the [Goddard Institute for Space Studies](#) (GISS), USA, in published anomaly values for the months January 1910 and January 2000.

The administrative upsurge of the temperature increase from January 1915 to January 2000 has grown from 0.45 (reported May 2008) to 0.67°C (reported June 2020). This represents an about 49% administrative temperature increase over this period, meaning that about half of the apparent global temperature increase from January 1910 to January 2000 (as reported by GISS) is due to administrative changes of the original data since May 2008.

Comparing global surface air temperature and lower troposphere satellite temperatures; updated to April 2020



10

Plot showing the average of monthly global surface air temperature estimates (HadCRUT4, GISS and NCDC) and satellite-based temperature estimates (RSS MSU and UAH MSU). The thin lines indicate the monthly value, while the thick lines represent the simple running 37-month average, nearly corresponding to a running 3-yr average. The lower panel shows the monthly difference between average surface air temperature and satellite temperatures. As the base period differs for the different temperature estimates, they have all been normalised by comparing to the average value of 30 years from January 1979 to December 2008.

Global air temperature linear trends updated to April 2020




Diagram showing the latest 5, 10, 20 and 30-yr linear annual global temperature trend, calculated as the slope of the linear regression line through the data points, for two satellite-based temperature estimates (UAH MSU and RSS MSU).

11






Diagram showing the latest 5, 10, 20, 30, 50, 70 and 100-year linear annual global temperature trend, calculated as the slope of the linear regression line through the data points, for three surface-based temperature estimates (GISS, NCDC and HadCRUT4).

All in one, Quality Class 1, 2 and 3; updated to April 2020




Superimposed plot of Quality Class 1 (UAH and RSS) global monthly temperature estimates. As the base period differs for the individual temperature estimates, they have all been normalised by comparing with the average value of the initial 120 months (30 years) from January 1979 to December 2008. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of both temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the individual 1979-2008 averages.

12



Superimposed plot of Quality Class 1 and 2 (UAH, RSS and HadCRUT4) global monthly temperature estimates. As the base period differs for the individual temperature estimates, they have all been normalised by comparing with the average value of the initial 120 months (30 years) from January 1979 to December 2008. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of all three temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the individual 1979-2008 averages.



Superimposed plot of Quality Class 1, 2 and 3 global monthly temperature estimates (UAH, RSS, HadCRUT4, GISS and NCDC). As the base period differs for the individual temperature estimates, they have all been normalised by comparing with the average value of the initial 120 months (30 years) from January 1979 to December 2008. The heavy black line represents the simple running 37 month (c. 3 year) mean of the average of all five temperature records. The numbers shown in the lower right corner represent the temperature anomaly relative to the individual 1979-2008 averages.

Please see notes on page 8 relating to the above three quality classes.

Satellite- and surface-based temperature estimates are derived from different types of measurements, and that comparing them directly as done in the diagrams above therefore may be questionable.


However, as both types of estimate often are discussed together, the above composite diagrams may nevertheless be of interest. In fact, the different types of temperature estimates appear to agree as to the overall temperature variations on a 2-3-year scale, although on a shorter time scale there are often considerable differences between the individual records. However, since about 2003 the surface records are slowly drifting towards higher temperatures than the combined satellite record (see p. 10), although this difference recently was much reduced by the adjustment of the RSS satellite series (see lower diagram on page 5).

The combined records (diagram on p. 13) suggest a modest global air temperature increase since 1998,

about 0.15°C per decade. The year 1998 was, however, affected by a strong oceanographic El Niño event. Likewise, the recent (2015-16) strong El Niño event probably also represents a relatively short-lived spike on a longer development. It should also be noted that the apparent temperature increase since about 2003 at least partly is the result of ongoing administrative adjustments (page 5-9). At the same time, the temperature records considered in the present report do not indicate any temperature decrease over the last 20 years. See also diagram on page 48.

The present temperature development does not exclude the possibility that global temperatures may begin to increase significantly later. On the other hand, it also remains a possibility that Earth just now is passing an overall temperature peak, and that global temperatures may begin to decrease during the coming years. Again, time will show which of these possibilities is correct.

Global sea surface temperature, updated to May 2020



14

Sea surface temperature anomaly on 23 May 2020. Map source: Plymouth State Weather Center. Reference period: 1977-1991.


Because of the large surface areas near Equator, the temperature of the surface water in these regions is especially important for the global atmospheric temperature (p. 5-7). In fact, no less than 50% of planet Earth's surface area is located within 30°N and 30°S.

A mixture of relatively warm and cold water dominates much of the oceans, but with notable differences from month to month. All such ocean surface temperature changes will be influencing global air temperatures in the months to come.

The significance of any short-term cooling or warming reflected in air temperatures should not be


overstated. Whenever Earth experiences cold La Niña or warm El Niño episodes (Pacific Ocean) major heat exchanges take place between the Pacific Ocean and the atmosphere above, sooner or later showing up in estimates of the global air temperature.

However, this does not necessarily reflect similar changes in the total heat content of the atmosphere-ocean system. In fact, global net changes can be small and such heat exchanges may mainly reflect redistribution of energy between ocean and atmosphere. What matters is the overall temperature development when seen over several years.




Global monthly average lower troposphere temperature over oceans (thin line) since 1979 according to [University of Alabama](#) at Huntsville, USA. The thick line is the simple running 37-month average. Insert: Argo global ocean temperature anomaly from floats, displaced vertically to make visual comparison easier.

15




Global monthly average sea surface temperature since 1979 according to University of East Anglia's [Climatic Research Unit \(CRU\)](#), UK. Base period: 1961-1990. The thick line is the simple running 37-month average. Insert: Argo global ocean temperature anomaly from floats, displaced vertically to make visual comparison easier.



Global monthly average sea surface temperature since 1979 according to the [National Climatic Data Center](#) (NCDC), USA. Base period: 1901-2000. The thick line is the simple running 37-month average. Insert: Argo global ocean temperature anomaly from floats, displaced vertically to make visual comparison easier.


June 18, 2015: NCDC has introduced several rather large administrative changes to their sea surface temperature record. The overall result is to produce a record giving the impression of a continuous temperature increase, also in the 21st century. As the oceans cover about 71% of the entire surface of planet Earth, the effect of this adjustment is clearly reflected in the NCDC record for global surface air temperature (p. 7).

Ocean temperature in uppermost 100 m, updated to March 2020




World Oceans vertical average temperature 0-100 m depth since 1955. The thin line indicates 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](#). Base period 1955-2010.


17



Pacific Ocean vertical average temperature 0-100 m depth since 1955. The thin line indicates 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](#). Base period 1955-2010.




Atlantic Ocean vertical average temperature 0-100 m depth since 1955. The thin line indicates 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](#). Base period 1955-2010.




Indian Ocean vertical average temperature 0-100 m depth since 1955. The thin line indicates 3-month values, and the thick line represents the simple running 39-month (c. 3 year) average. Data source: [NOAA National Oceanographic Data Center \(NODC\)](#). Base period 1955-2010.

North Atlantic heat content uppermost 700 m, updated to March 2020




19




Global monthly heat content anomaly (10^{18} Joules) in the uppermost 700 m of the North Atlantic (60-0W, 30-65N; see map above) ocean since January 1955. The thin line indicates monthly values, and the thick line represents the simple running 37-month (c. 3 year) average. Data source: [National Oceanographic Data Center](#) (NODC).

North Atlantic temperatures 0-800 m depth along 59°N, 30-0W, updated to July 2019




Time series depth-temperature diagram along 59 N across the North Atlantic Current from 30°W to 0°W, from surface to 800 m depth. Source: [Global Marine Argo Atlas](#). See also the diagram below.

20



Average temperature along 59 N, 30-0W, 0-800m depth, corresponding to the main part of the North Atlantic Current, using [Argo](#)-data. Source: [Global Marine Argo Atlas](#). Additional information can be found in: Roemmich, D. and J. Gilson, 2009. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. [Progress in Oceanography](#), 82, 81-100.

Global ocean temperature 0-1900 m depth summary, updated to July 2019




Summary of average temperature in uppermost 1900 m in different parts of the global oceans, using [Argo](#)-data. Source: [Global Marine Argo Atlas](#). Additional information can be found in: Roemmich, D. and J. Gilson, 2009. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. [Progress in Oceanography](#), 82, 81-100.

The global summary diagram above shows that, on average, the temperature of the global oceans down to 1900 m depth has been increasing since about 2011. It is also seen that this increase since 2013 dominantly is due to oceanic changes occurring near the Equator, between


30°N and 30°S. In contrast, for the circum-Arctic oceans north of 55°N, depth-integrated ocean temperatures have been decreasing since 2011. Near the Antarctic, south of 55°S, temperatures have essentially been stable. At most latitudes, a clear annual rhythm is seen.

Global ocean net temperature change since 2004 at different depths, updated to July 2019



Net temperature change since 2004 from surface to 1900 m depth in different parts of the global oceans, using [Argo](#)-data. Source: [Global Marine Argo Atlas](#). Additional information can be found in: Roemmich, D. and J. Gilson, 2009. The 2004-2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program. [Progress in Oceanography](#), 82, 81-100. Please note that due to the spherical form of Earth, northern and southern latitudes represent only small ocean volumes, compared to latitudes near the Equator.

La Niña and El Niño episodes, updated to May 2020




23

Warm ($>+0.5^{\circ}\text{C}$) and cold ($<0.5^{\circ}\text{C}$) episodes for the [Oceanic Niño Index \(ONI\)](#), defined as 3 month running mean of ERSSTv4 SST anomalies in the Niño 3.4 region (5°N - 5°S , 120° - 170°W). For historical purposes cold and warm episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons. Anomalies are centred on 30-yr base periods updated every 5 years.

The recent 2015-16 El Niño episode is among the strongest since the beginning of the record in 1950. Considering the entire record, however, recent


variations between El Niño and La Niña episodes do not appear abnormal in any way.

Troposphere and stratosphere temperatures from satellites, updated to May 2020



Global monthly average temperature in different according to University of Alabama at Huntsville, USA. The thin lines represent the monthly average, and the thick line the simple running 37-month average, nearly corresponding to a running 3-year average.

Zonal lower troposphere temperatures from satellites, updated to May 2020




25

Global monthly average lower troposphere temperature since 1979 for the tropics and the northern and southern extratropics, according to University of Alabama at Huntsville, USA. Thin lines show the monthly temperature. Thick lines represent the simple running 37-month average, nearly corresponding to a running 3-year average. Reference period 1981-2010.

The overall warming since 1980 has dominantly been a northern hemisphere phenomenon, and mainly played out as a marked change between 1994 and 1999. However, this rather rapid temperature change is influenced by the Mt. Pinatubo eruption 1992-93 and the

subsequent 1997 El Niño episode. The diagram also shows the temperature effects of the strong Equatorial El Niño's in 1997 and 2015-16, as well as the moderate El Niño in 2019, apparently were spreading to higher latitudes in both hemispheres with some delay.

Arctic and Antarctic lower troposphere temperature, updated to May 2020



Global monthly average lower troposphere temperature since 1979 for the North Pole and South Pole regions, based on satellite observations ([University of Alabama](#) at Huntsville, USA). Thin lines show the monthly temperature. The thick line is the simple running 37-month average, nearly corresponding to a running 3-year average. Reference period 1981-2010.


In the Arctic region, warming mainly took place 1994-96, and less so subsequently. In 2016, however, temperatures peaked for several months, presumably because of oceanic heat given off to the atmosphere during the recent El Niño 2015-16 (see also figure on page 23) and then advected to higher latitudes.

This underscores how Arctic air temperatures may be affected not only by variations in local conditions but also by variations playing out in geographically remote

regions. An overall temperature decrease has characterised the Arctic since 2016 (see also diagrams on page 28-30).

In the Antarctic region, temperatures have remained almost stable since the onset of the satellite record in 1979. In 2016-17 a small temperature peak visible in the monthly record may be interpreted as the subdued effect of the recent El Niño episode.

Temperature over land versus over oceans, updated to May 2020



Global monthly average lower troposphere temperature since 1979 measured over land and oceans, respectively, according to [University of Alabama](#) at Huntsville, USA. Thick lines are the simple running 37-month average, nearly corresponding to a running 3-year average. Reference period 1981-2010.

Since 1979, the lower troposphere over land has warmed much more than over oceans, suggesting that the overall warming is derived mainly from incoming solar radiation.

In addition, there may be other reasons for this divergence, such as, e.g., variations in cloud cover and land use.

Arctic and Antarctic surface air temperature, updated to April 2020




Diagram showing area weighted Arctic (70-90°N) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 2000, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.

28




Diagram showing area weighted Antarctic (70-90°S) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 2000, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.




Diagram showing area weighted Arctic (70-90°N) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 1957, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.




Diagram showing area weighted Antarctic (70-90°S) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 1957, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.




Diagram showing area-weighted Arctic (70-90°N) monthly surface air temperature anomalies ([HadCRUT4](#)) since January 1920, in relation to the WMO [normal period](#) 1961-1990. The thin line shows the monthly temperature anomaly, while the thicker line shows the running 37-month (c. 3 year) average.

Because of the relatively small number of Arctic stations before 1930, month-to-month variations in the early part of the Arctic temperature record 1920-2018 are bigger than later (diagram above).

The period from about 1930 saw the establishment of many new Arctic meteorological stations, first in Russia and Siberia, and following the 2nd World War, also in North America. The period since 2005 is warm, about as warm as the period 1930-1940.


As the HadCRUT4 data series has improved high latitude coverage data coverage (compared to the HadCRUT3 series), the individual 5°x5° grid cells have been weighted according to their surface area. This area correction is

especially important for polar regions. This approach differs from the approach used by Gillett et al. 2008, which calculated a simple average, without any correction for the substantial surface area effect of latitude in polar regions.

Literature:


Gillett, N.P., Stone, D.A., Stott, P.A., Nozawa, T., Karpechko, A.Y.U., Hegerl, G.C., Wehner, M.F. and Jones, P.D. 2008. Attribution of polar warming to human influence. *Nature Geoscience* 1, 750-754.

Arctic and Antarctic sea ice, updated to May 2020




31

Sea ice extent 23 May 2020. The median limit of sea ice (orange line) is defined as 15% sea ice cover, according to the average of satellite observations 1981-2010 (both years included). Sea ice may therefore well be encountered outside and open water areas inside the limit shown in the diagrams above. Map source: National Snow and Ice Data Center (NSIDC).



Diagrams showing Arctic sea ice extent and concentration 24 May 2019 (left) and 2020 (right), according to the Japan Aerospace Exploration Agency (JAXA).



Graphs showing monthly Antarctic, Arctic and global sea ice extent since November 1978, according to the [National Snow and Ice data Center \(NSIDC\)](#).

32






Diagram showing daily Arctic sea ice extent since June 2002, to 24 May 2020, by courtesy of [Japan Aerospace Exploration Agency \(JAXA\)](#).


Sea Ice Thickness, 24-May-2020



33



Diagrams showing Arctic sea ice extent and thickness 24 May 2019 (left) and 2020 (right and above) and the seasonal cycles of the calculated total arctic sea ice volume, according to [The Danish Meteorological Institute \(DMI\)](#). The mean sea ice volume and standard deviation for the period 2004-2013 are shown by grey shading.



12 month running average sea ice extension, global and in both hemispheres since 1979, the satellite-era. The October 1979 value represents the monthly 12-month average of November 1978 - October 1979, the November 1979 value represents the average of December 1978 - November 1979, etc. The stippled lines represent a 61-month (ca. 5 years) average. Data source: National Snow and Ice Data Center (NSIDC).

Sea level in general

Global (or eustatic) sea-level change is measured relative to an idealised reference level, the geoid, which is a mathematical model of planet Earth's surface (Carter et al. 2014). Global sea-level is a function of the volume of the ocean basins and the volume of water they contain. Changes in global sea-level are caused by – but not limited to - four main mechanisms:

1. Changes in local and regional air pressure and wind, and tidal changes introduced by the Moon.
2. Changes in ocean basin volume by tectonic (geological) forces.
3. Changes in ocean water density caused by variations in currents, water temperature and salinity.
4. Changes in the volume of water caused by changes in the mass balance of terrestrial glaciers.

In addition to these there are other mechanisms influencing sea-level; such as storage of ground water, storage in lakes and rivers, evaporation, etc.

Mechanism 1 is controlling sea-level at many sites on a time scale from months to several years. As an example, many coastal stations show a pronounced annual variation reflecting seasonal changes in air pressures and wind speed. Longer-term climatic changes playing out over decades or centuries will also affect measurements of sea-level changes. Hansen et al. (2011, 2015) provide excellent analyses of sea-level changes caused by recurrent changes of the orbit of the Moon and other phenomena.

Mechanism 2 – with the important exception of earthquakes and tsunamis - typically operates over long (geological) time scales and is not significant on human time scales. It may relate to variations in the seafloor spreading rate, causing volume changes in mid-ocean mountain ridges, and to the slowly changing configuration of land and oceans. Another effect may be the slow rise of basins due to isostatic offloading by deglaciation after an ice age. The floor of the Baltic Sea and the Hudson Bay are presently rising, causing a slow net transfer of

water from these basins into the adjoining oceans. Slow changes of very big glaciers (ice sheets) and movements in the mantle will affect the gravity field and thereby the vertical position of the ocean surface. Any increase of the total water mass as well as sediment deposition into oceans increase the load on their bottom, generating sinking by viscoelastic flow in the mantle below. The mantle flow is directed towards the surrounding land areas, which will rise, thereby partly compensating for the initial sea level increase induced by the increased water mass in the ocean.

Mechanism 3 (temperature-driven expansion) only affects the uppermost part of the oceans on human time scales. Usually, temperature-driven changes in density are more important than salinity-driven changes. Seawater is characterised by a relatively small coefficient of expansion, but the effect should however not be overlooked, especially when interpreting satellite altimetry data. Temperature-driven expansion of a column of seawater will not affect the total mass of water within the column considered and will therefore not affect the potential at the top of the water column. Temperature-driven ocean water expansion will therefore not in itself lead to any lateral displacement of water, but only locally lift the ocean surface. Near the coast, where people are living, the depth of water approaches zero, so no measurable temperature-driven expansion will take place here (Mörner 2015). Mechanism 3 is for that reason not important for coastal regions.


Mechanism 4 (changes in glacier mass balance) is an important driver for global sea-level changes along coasts, for human time scales. Volume changes of floating glaciers – ice shelves – has no influence on the global sea-level, just like volume changes of floating sea ice has no influence. Only the mass-balance of grounded or land-based glaciers is important for the global sea-level along coasts.

Summing up: Presumably, mechanism 1 and 4 are the most important for understanding sea-level changes along coasts.

References:

- Carter R.M., de Lange W., Hansen, J.M., Humlum O., Idso C., Kear, D., Legates, D., Mörner, N.A., Ollier C., Singer F. & Soon W. 2014. Commentary and Analysis on the Whitehead& Associates 2014 NSW Sea-Level Report. Policy Brief, NIPCC, 24. September 2014, 44 pp. <http://climatechangereconsidered.org/wp-content/uploads/2014/09/NIPCC-Report-on-NSW-Coastal-SL-9z-corrected.pdf>
- Hansen, J.-M., Aagaard, T. and Binderup, M. 2011. Absolute sea levels and isostatic changes of the eastern North Sea to central Baltic region during the last 900 years. *Boreas*, 10.1111/j.1502-3885.2011.00229.x. ISSN 0300-9483.
- Hansen, J.-M., Aagaard, T. and Huijpers, A. 2015. Sea-Level Forcing by Synchronization of 56- and 74-Year Oscillations with the Moon's Nodal Tide on the Northwest European Shelf (Eastern North Sea to Central Baltic Sea). *Journ. Coastal Research*, 16 pp.
- Mörner, Nils-Axel 2015. Sea Level Changes as recorded in nature itself. *Journal of Engineering Research and Applications*, Vol.5, 1, 124-129.

Global sea level from satellite altimetry, updated to January 2018



Global sea level since December 1992 according to the Colorado Center for Astrodynamics Research at University of Colorado at Boulder. The blue dots are the individual observations, and the purple line represents the running 121-month (ca. 10 year) average. The two lower panels show the annual sea level change, calculated for 1 and 10-year time windows, respectively. These values are plotted at the end of the interval considered. Data from the TOPEX/Poseidon mission have been used before 2002, and data from the Jason-1 mission (satellite launched December 2001) after 2002.


Ground truth is a term used in various fields to refer to information provided by direct observation as opposed to information provided by inference, such as, e.g., by satellite observations.

In remote sensing using satellite observations, ground truth data refers to information collected on location. Ground truth allows the satellite data to be related to real features observed on the planet surface. The collection of ground truth data enables calibration of remote-sensing

data, and aids in the interpretation and analysis of what is being sensed or recorded by satellites. Ground truth sites allow the remote sensor operator to correct and improve the interpretation of satellite data.

For satellite observations on sea level ground true data are provided by the classical tide gauges (example diagram on next page), that directly measures the local sea level many places distributed along the coastlines on the surface of the planet.

Global sea level from tide-gauges, updated to December 2018



Holgate-9 monthly tide gauge data from PSMSL Data Explorer. *Holgate (2007)* suggested the nine stations listed in the diagram to capture the variability found in a larger number of stations over the last half century studied previously. For that reason, average values of the *Holgate-9* group of tide gauge stations are interesting to follow, even though Auckland (New Zealand) has not reported data since 2000, and Cascais (Portugal) not since 1993. Unfortunately, by this data loss the *Holgate-9* series since 2000 is underrepresented with respect to the southern hemisphere. The blue dots are the individual average monthly observations, and the purple line represents the running 121-month (ca. 10 year) average. The two lower panels show the annual sea level change, calculated for 1 and 10-year windows, respectively. These values are plotted at the end of the interval considered.

Data from tide-gauges all over the world suggest an average global sea-level rise of 1-1.5 mm/year, while the satellite-derived record (page 36) suggest a rise of about 3.2 mm/year, or more. The noticeable difference (at least 1:2) between the two data sets is remarkable but has no


broadly accepted explanation. It is however known that satellite observations are facing several complications in areas near the coast. Vignudelli et al. (2019) provide an updated overview of the current limitations of classical satellite altimetry in coastal regions.

References:

Holgate, S.J. 2007. On the decadal rates of sea level change during the twentieth century. *Geophys. Res. Letters*, 34, L01602, doi:10.1029/2006GL028492


Vignudelli et al. 2019. Satellite Altimetry Measurements of Sea Level in the Coastal Zone. *Surveys in Geophysics*, Vol. 40, p. 1319–1349. <https://link.springer.com/article/10.1007/s10712-019-09569-1>

Northern Hemisphere weekly and seasonal snow cover, updated to May 2020




Northern hemisphere snow cover (white) and sea ice (yellow) 24 May 2019 (left) and 2020 (right). Map source: [National Ice Center \(NIC\)](#).


38



Northern hemisphere weekly snow cover since January 2000 according to Rutgers University Global Snow Laboratory. The thin blue line is the weekly data, and the thick blue line is the running 53-week average (approximately 1 year). The horizontal red line is the 1972-2019 average.




Northern hemisphere weekly snow cover since January 1972 according to Rutgers University Global Snow Laboratory. The thin blue line is the weekly data, and the thick blue line is the running 53-week average (approximately 1 year). The horizontal red line is the 1972-2019 average.



Northern hemisphere seasonal snow cover since January 1972 according to Rutgers University Global Snow Laboratory.

Atmospheric specific humidity, updated to May 2020



[Specific atmospheric humidity](#) (g/kg) at three different altitudes in the lower part of the atmosphere ([the Troposphere](#)) since January 1948 ([Kalnay et al. 1996](#)). The thin blue lines show monthly values, while the thick blue lines show the running 37-month average (about 3 years). Data source: [Earth System Research Laboratory \(NOAA\)](#).


Water vapor is the most important greenhouse gas in the Troposphere. The highest concentration is found within a latitudinal range from 50°N to 60°S. The two polar regions of the Troposphere are comparatively dry.

The diagram above shows the specific atmospheric humidity to be stable or slightly increasing up to about 4-5 km altitude. At higher levels in the Troposphere (about 9 km), the specific humidity has been decreasing for the duration of the record (since 1948), but with shorter

variations superimposed on the falling trend. A Fourier frequency analysis (not shown here) shows these variations to be influenced especially by a periodic variation of about 3.7-year duration.


The persistent decrease in specific humidity at about 9 km altitude is particularly noteworthy, as this altitude roughly corresponds to the level where the theoretical temperature effect of increased atmospheric CO₂ is expected initially to play out.

Atmospheric CO₂, updated to May 2020



Monthly amount of atmospheric CO₂ (upper diagram) and annual growth rate (lower diagram); average last 12 months minus average preceding 12 months, thin line) of atmospheric CO₂ since 1959, according to data provided by the [Mauna Loa Observatory](#), Hawaii, USA. The thick, stippled line is the simple running 37-observation average, nearly corresponding to a running 3-year average

The relation between annual change of atmospheric CO₂ and La Niña and El Niño episodes, updated to May 2020



Visual association between annual growth rate of atmospheric CO₂ (upper panel) and Oceanic Niño Index (lower panel). See also diagrams on page 40 and 22, respectively.

Changes in the global atmospheric CO₂ is seen to vary roughly in concert with changes in the Oceanic Niño Index. The typical sequence of events is that changes in the global atmospheric CO₂ to a certain degree follows changes in the Oceanic Niño Index, but clearly not in all details. Many processes, natural as well as human, controls the amount of atmospheric CO₂, but oceanographic processes are clearly important (see also diagram on next page).

Atmospheric CO₂ and the present coronavirus pandemic


Modern political initiatives usually assume the human influence (mainly the burning of fossil fuels) to represent

the core reason for the observed increase in atmospheric CO₂ since 1958 (see diagrams on page 41).

The present (since January 2020) coronavirus pandemic has resulted in a marked reduction in the global consumption of fossil fuels, as is well reflected by plummeting value of oil and gas. It is interesting to follow the effect of this on the amount of atmospheric CO₂.

By the end of May 2020 there is still no clear effect to be seen. The simplest explanation for this is that the human contribution is too small compared to the numerous natural sources and sinks for atmospheric CO₂ to appear in diagrams showing the amount of atmospheric CO₂ (diagrams on p. 41-43).

The phase relation between atmospheric CO₂ and global temperature, updated to April 2020



12-month change of global atmospheric CO₂ concentration ([Mauna Loa](#); green), global sea surface temperature ([HadSST3](#); blue) and global surface air temperature ([HadCRUT4](#); red dotted). All graphs are showing monthly values of DIFF12, the difference between the average of the last 12 months and the average for the previous 12 months for each data series.

43

The typical sequence of events is seen to be that changes in the global atmospheric CO₂ follow changes in global surface air temperature, which again follow changes in global ocean surface temperatures. Thus, changes in


global atmospheric CO₂ are lagging 9.5–10 months behind changes in global air surface temperature, and 11–12 months behind changes in global sea surface temperature.

References:


Humlum, O., Stordahl, K. and Solheim, J-E. 2012. The phase relation between atmospheric carbon dioxide and global temperature. *Global and Planetary Change*, August 30, 2012.


<http://www.sciencedirect.com/science/article/pii/S0921818112001658?v=s5>

Global air temperature and atmospheric CO₂, updated to May 2020





44





45





Diagrams showing UAH, RSS, HadCRUT4, NCDC and GISS monthly global air temperature estimates (blue) and the monthly atmospheric CO₂ content (red) according to the [Mauna Loa Observatory](#), Hawaii. The Mauna Loa data series begins in March 1958, and 1958 was therefore chosen as starting year for the all diagrams above. Reconstructions of past atmospheric CO₂ concentrations (before 1958) are not incorporated in this diagram, as such past CO₂ values are derived by other means (ice cores, stomata, or older measurements using different methodology), and therefore are not directly comparable with direct atmospheric measurements. The dotted grey line indicates the approximate linear temperature trend, and the boxes in the lower part of the diagram indicate the relation between atmospheric CO₂ and global surface air temperature, negative or positive.

Most climate models are programmed to give the greenhouse gas carbon dioxide CO₂ significant influence on the modelled global temperature. It is therefore relevant to compare different temperature records with measurements of atmospheric CO₂, as shown in the diagrams above.

Any comparison, however, should not be made on a monthly or annual basis, but for a longer time, as other effects (oceanographic, cloud cover, etc.) may override the potential influence of CO₂ on short time scales such as just a few years.

It is of cause equally inappropriate to present new meteorological record values, whether daily, monthly or

annual, as demonstrating the legitimacy of the hypothesis ascribing high importance of atmospheric CO₂ for global temperatures. Any such meteorological record value may well be the result of other phenomena. Unfortunately, many news media repeatedly fall into this trap.

What exactly defines the critical length of a relevant period length to consider for evaluating the alleged importance of CO₂ remains elusive and represents a theme for discussion. However, the length of the critical period must be inversely proportional to the temperature sensitivity of CO₂, including feedback effects. Thus, if the net temperature effect of atmospheric CO₂ is strong, the critical period will be short, and vice versa.


However, past climate research history provides some clues as to what has traditionally been considered the relevant length of period over which to compare temperature and atmospheric CO₂. After about 10 years of concurrent global temperature- and CO₂-increase, IPCC was established in 1988. For obtaining public and political support for the CO₂-hypothesis the 10-year warming period leading up to 1988 most likely was considered important. Had the global temperature instead been decreasing at that time, political support for the hypothesis would have been difficult to obtain in 1988.

Based on the previous 10 years of concurrent temperature- and CO₂-increase, many climate

scientists in 1988 presumably felt that their understanding of climate dynamics was enough to conclude about the importance of CO₂ for global temperature changes. From this it may safely be concluded that 10 years was considered a period long enough to demonstrate the effect of increasing atmospheric CO₂ on global temperatures. The 10-year period is also basis for the anomaly diagrams shown on page 2.

Adopting this approach as to critical time length (at least 10 years), the varying relation (positive or negative) between global temperature and atmospheric CO₂ has been indicated in the lower panels of the diagrams above.

Latest 20-year QC1 global monthly air temperature changes, updated to May 2020



48

Last 20 years' global monthly average air temperature according to Quality Class 1 (UAH and RSS; see p.10) global monthly temperature estimates. The thin blue line represents the monthly values. The thick black line is the linear fit, with 95% confidence intervals indicated by the two thin black lines. The thick green line represents a 5-degree polynomial fit, with 95% confidence intervals indicated by the two thin green lines. A few key statistics are given in the lower part of the diagram (please note that the linear trend is the monthly trend).

In the enduring scientific climate debate the following question is often put forward: Is the surface air temperature still increasing or has it basically remained without significant changes during the last 15-16 years?


The diagram above may be useful in this context and demonstrates the differences between two often used statistical approaches to determine recent temperature trends. Please also note that such fits only attempt to describe the past, and usually have small, if any, predictive power. In addition, before using any linear trend (or other) analysis of time series a proper statistical model should be chosen, based on statistical justification. For temperature time series, there is no *a priori* physical reason why the long-term trend should be linear in time.

In fact, climatic time series often have trends for which a straight line is not a good approximation, as is clearly demonstrated by several of the diagrams shown in the present report.

For an excellent description of problems often encountered by analyses of temperature time series analyses, please see [Keenan, D.J. 2014: Statistical Analyses of Surface Temperatures in the IPCC Fifth Assessment Report.](#)


See also diagrams on page 11.

Sunspot activity and QC1 average satellite global air temperature, updated to May 2020



Variation of global monthly air temperature according to Quality Class 1 (UAH and RSS; see p.4) and observed sunspot number as provided by the Solar Influences Data Analysis Center (SIDC), since 1979. The thin lines represent the monthly values, while the thick line is the simple running 37-month average, nearly corresponding to a running 3-year average. The asymmetrical temperature 'bump' around 1998 is influenced by the oceanographic El Niño phenomenon in 1998, as is the case also for 2015-16. Temperatures in year 2019 was influenced by a moderate El Niño.

1815: The Tambora volcanic eruption and the cold summer in New England 1816



The Central England temperature series 1770-1840. The length of the cooling effect of the Tambora 1815 eruption is indicated by the blue bar. These graphs has been prepared using the composite monthly meteorological series since 1659, originally painstakingly homogenized and published by the late professor Gordon Manley (1974). The data series is now updated by the Hadley Centre.

The April 1815 eruption of Tambora was probably the largest eruption in historic time. About 150 cubic kilometres of ash were erupted. This is about 150 times more than the 1980 eruption of Mount St. Helens in USA. Ash fell as far as 1,300 km from the volcano. In central Java and Kalimantan, 900 km from the eruption, one centimetre of ash fell. The eruption column is estimated to have reached a height of about 45 km, that is, reaching into the upper Stratosphere.

An estimated 92,000 people were killed by the 1815 Tambora eruption. About 10,000 direct deaths were caused by volcanic bomb impacts, tephra fall, and pyroclastic flows, the rest indirectly by starvation,

disease, and hunger. The eruption apparently lowered average world temperature by about 0.5-0.7°C over a period of 2-3 years. The 1815 eruption of Tambora was followed in North America and Europe by what was called "the year without a summer". Somewhat unusual, London experienced snow in August 1816.

In the years following the eruption air temperatures were low in many parts of the world, including New England in North America, where 1816 became known as the 'poverty year' (Perley 2001). In fact, the coldest summer known to have been experienced in New England was that of 1816. Many of the crops proved a failure and it seemed at the

time as if nothing would be produced. In New Hampshire but little pork was fattened on account of the scarcity and consequent great cost of corn, and the people used mackerel as a substitute for it.

There was frost and snow in all the summer months, and in the northwestern part of New England a severe drought prevailed, which added to the disastrous effects of the season.



51

Northeastern USA with New England as seen in Google Earth.

Perley (2001) states that in the fall of 1816 '*many people have endeavoured to ascertain some cause for the extraordinary nature of this summer, though no opinion has gained much ground. A large number of the people of that time believed that the large spots which appeared on the sun's disk that spring lessened the number of rays of light and consequently the earth was to that extent cooler than usual. The spots were so large that, for the first time in their history, they could be seen without the aid of telescope. The large sunspots were seen by the naked eye for several days, beginning on the third of May, and, reappearing on June 11, they were again seen for a few days only*'.

Clearly people also at that time (1816) were skilled observers of nature. However, the organized international communication of news was not

equally well developed. Apparently, few people, if any, in New England had received information on the Tambora eruption the previous year. Neither did they probably know that although changes in sunspot activity are capturing some important aspects of solar activity, periods with increasing sunspot activity usually signal increasing solar insolation, although it is still unclear how much of the solar variability is captured by changes in sunspot numbers. Clearly, sunspot activity is not the only important measure of solar activity. It is however likely that the at that time ongoing but mainly preceding Dalton sunspot minimum (1790-1820) in the year 1816 made people aware that a change towards higher sunspot activity was detectable on the Sun, and that this might represent a possible explanation for the unusual weather experienced.

Nevertheless, in New England May 1816 became both dry and unseasonably cold. At Chester, New Hampshire, newly ploughed land froze hard on 15 May, and snow fell in some of the northern parts of New England (Perley 2001). Near Weare, New Hampshire, there were no blossoms on the fruit trees until about 20 May. Throughout the entire summer the weather was the subject of conversations. People asked themselves and each other, if a change had not come over the climate, especially when they heard that in Ohio it had snowed on May 22, and on 13 May there had been frost as far south as Virginia (c.f. Perley 2001).

June 1816 began with some excessively hot days, but soon it became cold again (Perley 2001). At Chester, New Hampshire ice formed on ponds of standing water in the morning of 6 June, while snow fell in Vermont, New Hampshire, and Maine. The frost and cold chilled and killed the martins and other birds, and froze the ground, cutting down corn and potatoes. In Vermont, the snow melted as fast as it fell, but in Massachusetts it was blown about as in winter. At Waterbury, Vermont, new fallen snow lay 20-25 cm thick 8 June. The oldest inhabitants did not recollect such an extraordinary cold June as June 1816. Many sheep perished with the cold, birds flew into houses for shelter, and great numbers of them were found dead in the fields. Throughout Maine, vegetation seemed to have been suspended.

July 1816 was characterized by frost in northern New Hampshire, which did considerable injury to the crops, and in Amherst, New Hampshire, snow fell. On 8 July the frost was so severe at Franconia, New Hampshire, that it cut off all the beans (Perley 2001). Vegetables and fruits, however, was pleasantly free of blemish, as the cold weather had indeed annihilated the caterpillars and canker-worms, but the king-bird and others, which usually feed on such insects, now resorted for sustenance to cherry trees and pea vines.

In early August 1816 there again was frost in New England, and at Amherst snow fell (Perley 2001). Then the weather improved, becoming warm and pleasant. On 20 August cold weather however again returned. At Keene and Chester in New Hampshire frost killed a large part of the corn, potatoes, beans, and wine, and also injured many crops in Maine. The mountains in Vermont were covered by snow.

In September 1816 3-5 cm snow fell at Springfield, Massachusetts, and the Vermont mountains had then been covered with snow for several days (Perley 2001). At Hartford, Connecticut, it was as cold as it usually is in November. At Hallowell, Maine, frost killed the corn and injured potatoes in low grounds on 20 September. Before the end of September, snow fell at Boston for several hours.

Perley (2001) states the following about the general situation in New England following the cold and adverse weather during the summer 1816: *'There was great destitution among the people the next winter and spring. The farmers in some instances were reduced to the last extremity, and many cattle died. The poorer men could not buy corn at the exorbitant prices for which it was sold.In the autumn, stock was sold at extremely low prices on account of lack of hay and corn, a pair of four-year-old cattle being brought for thirty-nine dollars in Chester, New Hampshire.The next spring hay was sold in New Hampshire in a few instances as high as one hundred and eighty dollars per ton, its general price, however, being thirty dollars'*.

Small wonder that periods of cooling traditionally have been characterized as periods of '*climatic deterioration*', while periods of warming usually have been described as periods of '*climatic improvement*'.

References:

Perley, S. 2001. *Historic Storms of New England*. Commonwealth Editions, Beverly, Massachusetts, 302 pp. First published in 1891 by Salem Press Publishing and Printing Company, Salem, Massachusetts.

All diagrams in this report, along with any supplementary information, including links to data sources and previous issues of this newsletter, are freely available for download on www.climate4you.com

Yours sincerely,

Ole Humlum (Ole.Humlum@gmail.com)

Arctic Historical Evaluation and Research Organisation, Longyearbyen, Svalbard

20 June 2020.